

QUANTUM COHOMOLOGY OF GRASSMANNIANS MODULO SYMMETRIES

HARALD HENGELBROCK

ABSTRACT. The quantum cohomology of Grassmannians exhibits two symmetries related to the quantum product, namely a $\mathbb{Z}/n$ action and an involution related to complex conjugation. We construct a new ring by dividing out these symmetries in an ideal theoretic way and analyze its structure, which is shown to control the sum of all coefficients appearing in the product of cohomology classes. We derive a combinatorial formula for the sum of all Littlewood-Richardson coefficients appearing in the expansion of a product of two Schur polynomials.

1. INTRODUCTION

In the quantum Schubert calculus of the Grassmannians there appear two kinds of important symmetries, one a cyclic symmetry as described in [1], another is an involution related to complex conjugation, see [5, 9, 8]. In the representation of Schubert classes as complex functions on a finite set of points in complex space, these symmetries act by changing the “complex direction” of the function values. In order to understand the invariant aspects of quantum Schubert calculus it should therefore be rewarding to understand the absolute value of the Schubert class functions. It turns out that the value of Schubert classes on certain real points already determines an interesting numerical aspect of this structure. This leads to the definition of a quotient ring, which will be the coordinate ring of these points. Next, the quantum Schubert calculus modulo symmetries is already governed by the cup product of Schubert classes whose Young diagram is contained in a small subsquare. This is the content of Theorem 4.8.

We introduce a bilinear pairing on the quantum cohomology ring, which is just the sum over all coefficients in the Schubert basis expansion of a product of two classes. The ideal defining the quotient ring above is just the zero space of the bilinear pairing. In this sense, the quotient ring describes quantum Schubert calculus modulo “numerical equivalence”. This results in a simple combinatorial description of this pairing in the context of symmetric functions in Corollary 4.10.

Furthermore, we observe another symmetry related to a parameterization of points of the quantum cohomology spectrum by Young diagrams as in [8]. The double specialization formula of Stembridge, see [13], induces a strong symmetry of points and Schubert classes, which identifies two sets of structures on the quantum cohomology.

Date: November 4, 2002.

1991 Mathematics Subject Classification. 14N35, 14N15, 05E10.

Key words and phrases. Grassmannian, Quantum cohomology, Quantum Schubert calculus.

We wish to thank Alexander Postnikov for helpful remarks and conversations.

2. COMPLEX CONJUGATION AND $\mathbb{Z}/n$ ACTION

Notation 2.1. Consider $G(k, n)$, the Grassmannian of k -planes in $\mathbb{C}^n$ with tautological bundle S , and the quantum cohomology ring R with deformation parameter q set to 1. This ring has the following presentation, see [11]:

$$R = \mathbb{C}[\sigma_1, \dots, \sigma_k] / (Y_{n-k+1}, \dots, Y_n + (-1)^k) \\ l := n - k, \sigma_i := c_i(S), Y_i := s_i(S)$$

We will denote by S_λ the Schubert class associated to the Young diagram $\lambda = (\lambda_1, \dots, \lambda_l)$. Denote by G the set of Young diagrams contained in a (l, k) -rectangle. The ring R has a vector space basis consisting of Schubert classes represented by Young diagrams in G . The generators σ_i of R are represented by diagrams with i boxes in the first row and zero boxes in all others. Three point GW-invariants of cohomology classes will be denoted by $\langle A, B, C \rangle$ for general classes A, B, C . Poincaré duality on R will be denoted by the $\hat{}$ -symbol. We use “ $*$ ” for the quantum product, and “ $\cup$ ” for the classical cup product. For standard facts related to the quantum cohomology of Grassmannians we refer to [2].

On R , there is a $\mathbb{Z}/n$ action, given by multiplication with powers of σ_k , see [1, Section 7]. This action induces equalities

$$\langle \sigma_k^a * A, \sigma_k^b * B, \sigma_k^{n-a-b} C \rangle = \langle A, B, C \rangle$$

for $A, B, C \in R$, see [1, Proposition 7.2]. There is an involution on R , which is just complex conjugation, where we think of a cohomology class as a complex function on $\text{Spec } R$, see [5], or [9, 8], in a different context. This involution operates on Schubert classes by dualizing the subdiagrams right and below the Durfee square:




FIGURE 2.1. Complex conjugation

For $\lambda \sim (\lambda_1, \dots, \lambda_l)$, writing d_λ for the length of the Durfee square, $\overline{S_\lambda}$ will be represented by $(\mu_1, \dots, \mu_l)$ with

$$(2.1) \quad \mu_i = \begin{cases} d_\lambda + k - \lambda_{d_\lambda - i + 1} & : i \leq d_\lambda \\ d_\lambda - \lambda_{l - i + d_\lambda + 1} & : i > d_\lambda. \end{cases}$$

Algebraically, this involution can be written as

$$(2.2) \quad \overline{S_\lambda} = \widehat{S_\lambda * C_P},$$

where C_P is the class of a point on $G(k, n)$.

Complex conjugation yields an important equation of GW-invariants:

$$(2.3) \quad \langle A, B, C \rangle = \langle \widehat{A}, \widehat{B}, \overline{C} \rangle \quad A, B, C \in R.$$

This equation implies that complex conjugation is a partial inversion: $A * \overline{B} = \sum c_i C_i$, where c_i is the coefficient of A in $C_i * B$. By 2.3, for any class $A \in R$, multiplication by $A * \overline{A}$ induces a semipositive definite symmetric endomorphism of the vector space R . Denote by $P_1, \dots, P_N$ the points of $\text{Spec } R$, then A is

represented by the vector $\begin{pmatrix} A(P_1) \\ \vdots \\ A(P_N) \end{pmatrix}$, and $A * \overline{A}$ will be $\begin{pmatrix} |A(P_1)|^2 \\ \vdots \\ |A(P_N)|^2 \end{pmatrix}$.

Because of

$$\overline{\sigma_k^r * A} = \sigma_k^{n-r} * \overline{A},$$

the map $A \mapsto A * \overline{A}$ is invariant on the orbits of $\mathbb{Z}/n$ action and of course complex conjugation. The absolute value of Schubert classes should therefore encode its invariant aspects.

In the combinatorial setting, we have following property of the map

$$(2.4) \quad \varphi : A \mapsto A * \overline{A} :$$

we define for any $A = \sum c_\lambda S_\lambda \in R$

$$|A|^2 = \sum c_\lambda \overline{c_\lambda},$$

e.g. the square of the hermitian norm with respect to the Schubert basis. $|A|^2$ is the coefficient of 1 in the expansion of the product $A * \overline{A}$, since

$$\langle A, \overline{A}, C_P \rangle = \sum_{\lambda, \mu} c_\lambda \overline{c_\mu} \langle S_\lambda, \overline{S_\mu}, C_P \rangle = \sum_{\lambda} c_\lambda \overline{c_\lambda} \langle S_\lambda, \overline{S_\lambda}, C_P \rangle = \sum_{\lambda, \mu} c_\lambda \overline{c_\mu},$$

where we used 2.2. Since the quantum product is homogenous mod n , the quantum ring carries a natural “mod n ” grading. Define $R_0 \subseteq R$ to be the subring of homogenous elements of degree 0 mod n , and define R_{inv} to be the subring of classes invariant under complex conjugation, that is, the real classes. Then φ maps Schubert classes into the subring $R_0 \cap R_{inv}$. For Schubert classes S_λ, S_μ the value

$$|S_\lambda * S_\mu|^2 = S_\lambda * S_\mu * \overline{S_\lambda * S_\mu} |_1$$

depends only on the values of φ on S_λ, S_μ : The last expression is just the scalar product of the coefficient vectors of $\varphi(S_\lambda)$ and $\varphi(S_\mu)$. Below, we will see that a related map controls the sum $\sum c_\lambda$ in a similar way.

3. SYMMETRIES OF POINTS AND SCHUBERT CLASSES

There is the following description of the points of $\text{Spec } R$, see [8]. For any $\lambda \in G$ write $\lambda = (\lambda_1, \dots, \lambda_l)$. Consider the set I of all tuples $I_\lambda = (i_1, \dots, i_l)$ indexed by $\lambda \in G$, where $i_j = \frac{l+1}{2} + \lambda_j - j$. We fix ξ to be any primitive n -th root of unity, and, in case l even, $\xi^{\frac{1}{2}}$ to be any primitive $2n$ -th root of unity. Then to any I_λ we can associate a point $P_\lambda \in \text{Spec } R \hookrightarrow \mathbb{C}^k$ by the following map:

$$I_\lambda = (i_1, \dots, i_l) \mapsto (h_1(\xi^{i_1}, \dots, \xi^{i_l}), \dots, h_k(\xi^{i_1}, \dots, \xi^{i_l})),$$

where the h_i are the full homogenous symmetric polynomials of degree i in l variables. Abusing notation, we will sometimes write P_S for the point corresponding to any Schubert class S , so P_1 is the point associated to the empty diagram. $S_\lambda(P_1)$ is invariant under complex conjugation, therefore real, and also positive, see [8, Theorem 8.4].

Proposition 3.1. (*Double specialization formula*) *Consider the matrix $M := (S_\lambda(P_\mu))_{\lambda,\mu}$. M is symmetric modulo division by the P_1 , that is*

$$(3.1) \quad \frac{S_\lambda(P_\mu)}{S_\lambda(P_1)} = \frac{S_\mu(P_\lambda)}{S_\mu(P_1)}.$$

Proof. This identity is an easy consequence of the definition of Schur polynomials, see [12, Exercise 7.32a]:

As a symmetric function in k variables $x_1, \dots, x_k$, the Schur polynomial can be defined as

$$S_\lambda = \frac{\det \left(x_i^{\lambda_j + k - j} \right)_{1 \leq i, j \leq k}}{\det \left(x_i^{k - j} \right)_{1 \leq i, j \leq k}},$$

see [7], I (3.1). In this sense, $S_\lambda(P_\mu)$ will be $S_\lambda(\xi^{I_\mu})$. Proving the identity 3.1, because of cancellation, it is enough to calculate the left hand side with $I'_\mu = (\mu_1 + k - 1, \mu_2 + k - 2, \dots, \mu_k)$. By [8, Lemma 4.4], we can assume $l \leq k$. Filling up the Young diagrams with zero rows we can assume $l = k$. We need to prove:

$$\frac{S_\lambda(\xi^{I'_\mu})}{S_\lambda(\xi^{I'_1})} = \frac{S_\mu(\xi^{I'_\lambda})}{S_\mu(\xi^{I'_1})}.$$

$$\frac{S_\lambda(\xi^{I'_\mu})}{S_\lambda(\xi^{I'_1})} = \frac{\frac{\det \left(\xi^{\mu_i + k - i + \lambda_j + k - j} \right)_{i,j}}{\det \left(\xi^{\mu_i + k - i + k - j} \right)_{i,j}}}{\frac{\det \left(\xi^{k - i + \lambda_j + k - j} \right)_{i,j}}{\det \left(\xi^{k - j + k - j} \right)_{i,j}}} = \frac{\det \left(\xi^{\mu_i + k - i + \lambda_j + k - j} \right) \cdot \det \left(\xi^{k - i + k - j} \right)}{\det \left(\xi^{k - i + \lambda_j + k - j} \right) \cdot \det \left(\xi^{\mu_i + k - i + k - j} \right)}.$$

We observe that the expression on the right is symmetric in λ and μ . □

Remark 3.2. The identity 3.1 is a special case of a double specialization formula of Stembridge, see [13, Theorem 7.4]. This paper also includes reference to the Macdonald polynomial conjecture, which is a further generalization.

Example 3.3. $n = 4$, $k = 2$, $\xi = i$, $\xi^{\frac{1}{2}} = \frac{i+1}{\sqrt{2}}$:

	$S_{0,0}$	$S_{1,0}$	$S_{2,0}$	$S_{1,1}$	$S_{2,1}$	$S_{2,2}$
$P_{0,0}$	1	$\sqrt{2}$	1	1	$\sqrt{2}$	1
$P_{1,0}$	1	0	i	$-i$	0	-1
$P_{2,0}$	1	$\sqrt{2}i$	-1	-1	$-\sqrt{2}i$	1
$P_{1,1}$	1	$-\sqrt{2}i$	-1	-1	$\sqrt{2}i$	1
$P_{2,1}$	1	0	$-i$	i	0	-1
$P_{2,2}$	1	$-\sqrt{2}$	1	1	$-\sqrt{2}$	1

After division of the row corresponding to $P_{0,0}$ we get a symmetric matrix:

	$S_{0,0}$	$S_{1,0}$	$S_{2,0}$	$S_{1,1}$	$S_{2,1}$	$S_{2,2}$
$P_{0,0}$	1	1	1	1	1	1
$P_{1,0}$	1	0	i	$-i$	0	-1
$P_{2,0}$	1	i	-1	-1	$-i$	1
$P_{1,1}$	1	$-i$	-1	-1	i	1
$P_{2,1}$	1	0	$-i$	i	0	-1
$P_{2,2}$	1	-1	1	1	-1	1

We have the following result for the values of the S_λ on P_{σ_k} , or the value of σ_k on points P_λ :

Lemma 3.4. $\frac{S_\lambda(P_{\sigma_k})}{S_\lambda(P_1)} = \sigma_k(P_\lambda) = \xi^d$, where d is the degree of S_λ modulo n .

Proof. We note that, by definition, for any $I_\lambda = (i_1, \dots, i_l)$ the sum $\sum i_j$ is just the degree of S_λ . Now, taking e_i to be the i -th elementary symmetric polynomial and $\lambda^t = (\lambda_1^t, \dots, \lambda_k^t)$ the transposed Young diagram, $h_k(\xi^{I_\lambda}) = e_k(\xi^{I_{\lambda^t}})$, see [8, Proof of Lemma 4.4], and $e_k(\xi^{I_{\lambda^t}}) = \xi^{\lambda_1^t} \dots \xi^{\lambda_k^t} = \xi^{|I_{\lambda^t}|} = \xi^{|I_\lambda|}$. $\square$

One immediate consequence of this Lemma is that the number of orbits under the $\mathbb{Z}/n$ action is the same as the number of Schubert classes in R_0 , the subring of classes of degree zero modulo n :

Proposition 3.5. $\dim_{\mathbb{C}} R_0 = \#\{\text{Orbits under the } \mathbb{Z}/n \text{ action}\}$

Proof. The number of orbits under the $\mathbb{Z}/n$ action is equal to the dimension of $R/(\sigma_k - 1)$, since the relation $(\sigma_k - 1)$ identifies exactly Schubert classes in the same orbit. Since we are dividing out a hyperplane, $\dim R/(\sigma_k - 1)$ will be the number of points $P_\lambda \in \text{Spec } R$ with $\sigma_k(P_\lambda) = 1$. The statement now follows from the previous lemma. $\square$

The equivalent of the $\mathbb{Z}/n$ action on points is multiplication with roots of unity. Defining $\Xi = (\xi, \xi^2, \dots, \xi^k)$, the action is induced by component wise multiplication of the vector $P_\lambda \sim (\sigma_1(P_\lambda), \dots, \sigma_k(P_\lambda))$ with powers of Ξ .

4. QUANTUM COHOMOLOGY MODULO SYMMETRIES

Definition 4.1. (see Figure 4.1) For the cohomology ring G of the Grassmannian $G(k, n)$, with $l := n - k$, define the subvectorspace G_S to be spanned by the classes associated to Young diagrams contained in the $[\frac{k}{2}] \times [\frac{l}{2}]$ rectangle, where $[\]$ means rounding down. Abusing notation, we will sometimes consider G_S to be the set of Young diagrams contained in this rectangle.

We define a map from $G_S \times G_S \mapsto G$ by assigning two Schubert classes S_λ and S_μ the class $S_\lambda(S_\mu)$ defined by embedding the Young diagram of S_μ into the diagram of $\overline{S_\lambda}$ in the following way:

$$n = 8, k = 4, \lambda = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \mu = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}, \bar{\lambda} = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & & \square \\ \hline \end{array}$$

$$S_\lambda(S_\mu) \cong \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array}$$

Such an embedding is always possible, since for $S_\lambda \in G_S$, $\overline{S_\lambda}$ always has an empty $\begin{bmatrix} k \\ 2 \end{bmatrix} \times \begin{bmatrix} l \\ 2 \end{bmatrix}$ rectangle at the outer corner of the Durfee square. Numerically, with d_λ the length of the Durfee square, and $\lambda \cong (\lambda_1, \dots, \lambda_l)$, $\mu \cong (\mu_1, \dots, \mu_l)$, $S_\lambda(S_\mu)$ is given by $(\nu_1, \dots, \nu_l)$

$$(4.1) \quad \nu_i = \begin{cases} d_\lambda + k - \lambda_{d_\lambda - i + 1} & : i \leq d_\lambda \\ \mu_{i - d_\lambda} + d_\lambda - \lambda_{l - i + d_\lambda + 1} & : i > d_\lambda. \end{cases}$$




FIGURE 4.1. The $l \times k$ box with subrectangle G_S

Remark 4.2. The subsquare G_S is the biggest subrectangle with the property that quantum product is just ordinary cup product without any “cutting off”.

Proposition 4.3. *All Schubert classes of degree zero mod n invariant under complex conjugation are of type $T_\alpha(T_\alpha)$ for some $\alpha \in G_S$.*

Proof. Take a Schubert class $S_\lambda \in G_S$. We claim that the Young diagram $\overline{S_\lambda}(S_\lambda)$ is invariant. This follows from a direct calculation using 4.1 and 2.1.

Now let S_λ be a degree 0 invariant Schubert class. By 2.1,

$$\deg \overline{S_\lambda} = nd_\lambda - \deg S_\lambda,$$

so for S_λ we have $nd_\lambda = 2 \deg S_\lambda = 2na$ for some $a \in \mathbb{N}$. The length of the Durfee square is even, so we can reverse the construction. With $S_\lambda = (\lambda_1, \dots, \lambda_l)$, the corresponding class $S \in G_S$ is given by $\mu = (\mu_1, \dots, \mu_{\lfloor \frac{l}{2} \rfloor}, 0, \dots, 0)$,

$$\mu_i = \lambda_{d_\lambda/2+i} - d_\lambda/2, \quad i \leq \left\lfloor \frac{l}{2} \right\rfloor.$$

It follows from 2.1 that S_λ is already determined by the μ_i , and that μ is contained in G_S : the subdiagrams right and below of the Durfee square of μ correspond to lower halves of selfdual Young diagrams, so they can not extend over more than half of the length or height of the rectangle. Also, the Durfee square itself has length $\frac{d_\lambda}{2}$. $\square$

We prove the following important description for the product of two classes in G_S :

Proposition 4.4. *For $S_\lambda, S_\mu \in G_S$,*

$$(4.2) \quad \left\langle \overline{S_\lambda}, S_\mu, \widehat{T_\alpha(T_\beta)} \right\rangle = \sum_{T_\nu \in G_S} \left\langle \widehat{S_\lambda}, T_\alpha, T_\nu \right\rangle \left\langle T_\nu, T_\beta, \widehat{S_\mu} \right\rangle,$$

$$\text{and } \overline{S_\lambda} * S_\mu = \sum_{\alpha, \beta} \left\langle \overline{S_\lambda}, S_\mu, \widehat{T_\alpha(T_\beta)} \right\rangle T_\alpha(T_\beta).$$

Proof. We will use the description of quantum product by rim hook reduction as in [2], see also [3] for a simple proof. All diagrams with nontrivial reduction will have a structure as indicated in Figure 4.2, I.




FIGURE 4.2. Skew diagrams associated to $\overline{S}_\lambda \cup S_\mu$ and $\overline{S}_\lambda * S_\lambda$


We will now enumerate all semistandard skew tableaux appearing in $\overline{S}_\lambda \cup S_\mu$ whose associate word is a reverse lattice word and whose reduction is of shape $T_\alpha(T_\beta)$ for fixed α, β . Note that such diagrams must have β as middle subdiagram, since any removable rim hook must extend over the width and length of G_S , such that the reduction will again have a shape similar to Figure 4.2, with β inscribed in the middle. Now take the set of all diagrams reducing to $T_\alpha(T_\beta)$ by the rim hook reduction. For any such fixed diagram, the number of reverse lattice word completions is the same as for the skew diagram obtained by exchanging the position of the middle and left subdiagram as in Figure 4.2, II, by reason of commutativity and associativity of the product in the tableaux ring, see [4, §5.1]. It is easy to enumerate these skew tableaux: looking at the set of skew diagrams with content T_ν in the right and middle subdiagram we find that this number is just $\langle \overline{S}_\lambda, T_\nu, \widehat{T}_\alpha \rangle = \langle S_\lambda, \widehat{T}_\nu, \widehat{T}_\alpha \rangle = \langle \widehat{S}_\lambda, T_\nu, T_\alpha \rangle$, where we used 2.3, since the rim hook reduction is independent of the subdiagram inscribed in the middle. Now, for a fixed completion of the middle and right subdiagram of content T_ν , we can enumerate all possible completions of the left one. By one of the definitions of the Littlewood-Richardson rule, see [4, §5.2, Corollary 2], we find that this number is indeed $\langle T_\nu, T_\beta, \widehat{S}_\mu \rangle$. Summing up, we get 4.2. The argument above also shows that rim hook reduction will only result in diagrams of shape $T_\alpha(T_\beta)$ for some $\alpha, \beta \in G_S$. $\square$

Remark 4.5. Using 2.3, the formula 4.2 can be interpreted as a restricted associativity equation for the product $S_\lambda * \overline{T}_\alpha * T_\beta$. In this sense, the class $T_\alpha(T_\beta)$, which appears in $\overline{T}_\alpha * T_\beta$, corresponds to restricting the associativity equation to classes T_ν in G_S .

Example 4.6. Let $n = 12, k = 6, \lambda = \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}, \mu = \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}, \alpha = \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}, \beta = \begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$. The

coefficient of $T_\alpha(T_\beta) \cong \begin{smallmatrix} \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \end{smallmatrix}$ in $\overline{S}_\lambda * S_\mu$ is equal to the number of pairs of

semistandard skew tableaux on the shapes λ/α and μ/β of same content, satisfying the reverse lattice word condition, as in the figure below:



In our example, $\langle \overline{S_\lambda}, S_\mu, \widehat{T_\alpha(T_\beta)} \rangle$ will be 2.

Associate to $S_\lambda \in G_S$ the symmetric matrix

$$A_\lambda := \left(\langle \widehat{S_\lambda}, T_\alpha, T_\beta \rangle \right)_{\alpha, \beta \in G_S}.$$

The product $\overline{S_\lambda} * S_\mu$ is determined by the matrix product $A_\lambda * A_\mu^t$.

The strong combinatorial description of the product $\overline{S_\lambda} * S_\mu$ should be helpful for further understanding the “unlimited” cup product $S_\lambda * S_\mu = S_\lambda \cup S_\mu$. The relationship between $\overline{S_\lambda} * S_\mu$ and $S_\lambda * S_\mu$ reflects a symmetry between adding and removing skew diagrams in a quantum sense, see 2.3. In the following we will see that S_λ and $\overline{S_\lambda}$ display an identical “numerical” behaviour.

Definition 4.7. Define the class $P := \sum_\lambda S_\lambda$. The ideal $\text{Ann } P$ should be considered as consisting of relations $(S_\lambda - \phi(S_\lambda))$ where ϕ is a “symmetry map”. We will also consider the bilinear pairing $\langle A, B \rangle_s = \sum_{T_\lambda \in G} \langle A, B, T_\lambda \rangle = \langle A, B, P \rangle$, which is just the sum over all coefficients appearing in the Schubert basis representation of products of classes $A, B \in R$. $\text{Ann } P$ is obviously the zero space of $\langle \cdot, \cdot \rangle_s$ and can be considered to be the ideal of relations of numerically equivalent classes with respect to $\langle \cdot, \cdot \rangle_s$.

Taking $\sigma_0 := 1$, we wish to show the following

Theorem 4.8. *$\text{Ann } P \cong (\{\sigma_{k-i} - \sigma_i\}_i)$, and $R_P := R/\text{Ann } P$ as a vector space is isomorphic to G_S .*

Remark 4.9. We have $(\{\sigma_{k-i} - \sigma_i\}) = (\{\overline{\sigma_i} - \sigma_i\}, \sigma_k - 1) \supseteq \{S_\lambda - \overline{S_\lambda}, S_\lambda - \sigma_k^i S_\lambda : S_\lambda \in R\}$, which shows that $\text{Ann } P$ identifies all the orbits under the $\mathbb{Z}/n$ action and complex conjugation.

Let V be a vector space with basis $\{v_\lambda\}$ indexed by the Schubert classes in G_S . Then the map

$$(4.3) \quad \begin{aligned} \psi : G_S &\rightarrow V \\ S_\lambda &\mapsto \sum_{\nu \in G_S} \sum_{\lambda \in G_S} \langle \widehat{S}, T_\lambda, T_\mu \rangle v_\nu \end{aligned}$$

is an isomorphism of vector spaces, since the corresponding matrix can be arranged to be triangular with 1 on the diagonal: the coefficient of v_ν in $\psi(S_\lambda)$ will only be nonzero if the diagram ν is contained in λ , and 1 if $\nu = \lambda$. The bilinear form $\langle -, - \rangle_s$ simply becomes the standard scalar product under this map. Since, by the Theorem, $\langle -, - \rangle_s$ descends to R_P , we deduce the following Corollary, see Remark 4.2:

Corollary 4.10. *Consider the Littlewood-Richardson expansion of the product of two Schur polynomials in the ring of symmetric functions:*

$$S_\lambda * S_\mu = \sum c_{\lambda, \mu}^\nu S_\nu.$$

The sum $\sum_{\nu} c_{\lambda, \mu}^{\nu}$ is identical to

$$\sum_{\alpha, \beta, \nu} c_{\alpha, \nu}^{\lambda} \cdot c_{\beta, \nu}^{\mu}.$$

Proof. (of the Theorem) Define $I := (\{\sigma_{k-i} - \sigma_i\}_i)$. We first show that $I \subseteq \text{Ann } P$. It is obvious that $\sigma_k - 1 \in \text{Ann } P$. We will show $\overline{\sigma_i} - \sigma_i \in \text{Ann } P$, which by 4.4 implies $\sigma_{k-i} - \sigma_i \in \text{Ann } P$. By an application of complex conjugation we find $\sigma_i * P \mid \widehat{S_{\lambda}} = \overline{\sigma_i} * P \mid \widehat{S_{\lambda}}$:

$$\begin{aligned} \langle \sigma_i, P, S_{\lambda} \rangle &= \sum \langle \sigma_i, S_{\lambda}, T_i \rangle = \sum \langle \sigma_i, S_{\lambda}, T_i \rangle^2 = \sigma_i * S_{\lambda} * \overline{S_{\lambda}} \cup \widehat{\sigma_i} \\ &= \sum \langle \sigma_i, \overline{\sigma_i}, T_i \rangle \langle S_{\lambda}, \overline{S_{\lambda}}, T_i \rangle = \sum \langle \overline{\sigma_i}, S_{\lambda}, T_i \rangle^2 \\ &= \sum \langle \overline{\sigma_i}, S_{\lambda}, T_i \rangle = \langle \overline{\sigma_i}, P, S_{\lambda} \rangle, \end{aligned}$$

using 4.10, 4.4 and associativity of quantum multiplication together with the fact that $\langle \sigma_i, S_{\lambda}, T_i \rangle$ is either 1 or 0. We observe that the classes in G_S have no relations in $\text{Ann } P$: this can be translated into the fact that $\langle \cdot, \cdot \rangle_s = \langle \cdot, \cdot, P \rangle$ is nondegenerate on G_S , which follows from the discussion after the Theorem.

Now it is enough to show that the dimension of R/I does not exceed the number of Schubert classes in G_S . We observe that $\text{Spec } R/I$ consists of all real points $P_{\lambda} \in \text{Spec } R$ with $\sigma_k(P_{\lambda}) = 1$, since

$$(4.4) \quad \overline{\sigma_i} = \sigma_{k-i} * \sigma_k,$$

so the relations $\sigma_{k-i} - \sigma_i$ guarantee that the imaginary part of any P_{λ} with $\sigma_k(P_{\lambda}) = 1$ vanishes. But these points, by Proposition 3.1 and Lemma 3.4, correspond 1-1 to Schubert classes of degree zero mod n which are invariant under complex conjugation. By Proposition 4.3, these are parameterized by Schubert classes in G_S . We have proved

$$\#\{P_{\lambda} \in \text{Spec } R/I\} \leq \#\{S_{\lambda} \in G_S\},$$

which implies the result. $\square$

Remark 4.11. 1. The last statement in the previous proof can be seen directly by proving that every monomial in R can be represented mod I by monomials coming from classes in G_S . This is done by carrying out the identification of both generators $\sigma_i \sim \sigma_{k-i}$ and dual generators $\sigma^i \sim \sigma^{l-i}$.

2. The points of $\text{Spec } R_P$ are exactly the P_{λ} for which S_{λ} is invariant of degree zero mod n . R_P is the coordinate ring of these points and the quotient map $R \mapsto R_P$ is the restriction map. All elements of R_P are real functions.

3. It is obvious that all the Schubert classes in the orbit of one $S_{\lambda} \in G_S$ are mapped to this class under the composition

$$R \rightarrow R_P \cong G_S.$$

In general, a Schubert class will be mapped to a nonpositive linear combination of Schubert classes in G_S , as in the following example:

$$n = 8, k = 4 \Rightarrow G_S \sim G(2, 4).$$

Take

$$\lambda = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \square & & \\ \hline \end{array}.$$

In G_S , $[\lambda]$ is represented by

$$\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} - 2.$$

Under the isomorphism ψ of 4.3, $[\lambda]$ is represented by

$$\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + 3 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + 1.$$

Note that this is a positive linear combination.

5. SUMMARY AND OUTLOOK

In this paper we observed the following set of dualities:

$$\begin{aligned} \{\text{Orbits under } \mathbb{Z}/n \text{ action}\} &\approx R_0 \\ \{\text{Orbits under } \mathbb{Z}/n \text{ action and comp. conj.}\} &\approx R_0 \cap R_{inv} \\ R_P &\approx \{S_\lambda \in R_0 \cap R_{inv}\}. \end{aligned}$$

The symmetry of Proposition 3.1 hints at a stronger relation beyond the equality of dimensions as vector spaces. This is especially obvious in the third case, where, as in the case of R , we find a 1-1 correspondence between points of $\text{Spec } R_P$ and a ‘‘Schubert basis’’ of R_P . The ring R_P controls quantum Schubert calculus modulo numerical equivalence, which implies that the orbits under the actions behave numerically identical. We expect that the composition $R \rightarrow R_P \xrightarrow{\psi} V$ results in vectors with positive components for all Schubert classes. A positive combinatorial formula for this map would suggest a new rule for constructing the quantum product. This is especially interesting with regards to the positivity of the GW invariants, which is evident geometrically but not as clear algebraically.

It would be rewarding to find an explicit relationship between products $S_\lambda * S_\mu$ and $\overline{S}_\lambda * S_\mu$, since this would imply a better understanding of the classical cup product, e.g. with regards to the question as to what classes appear with nonzero coefficient. Also, there might be a natural generalization to the quantum product.

A part of the results for the Grassmannian should be extendable to other homogeneous varieties. For example, the variety of full flags in $\mathbb{C}^n$ carries a $\mathbb{Z}/n$ action, see [10], which acts by shifting the word of a permutation. Computations indicate that the zero space of $\langle \cdot, \cdot \rangle_s$ is generated by this action. In analogy to the Grassmannian, the quotient ring would have a vector space basis consisting e.g. of Schubert classes whose permutations fix n . The quantum product should be nondegenerate on this subset, that is, the linear map $\langle S_\omega, \cdot \rangle_s$ determines the Schubert class. In analogy to Proposition 3.5, the number of orbits, $(n-1)!$, is identical to the number of Schubert classes of degree zero mod n , because the number of permutations with t mod n inversions is indeed $(n-1)!$, see for example [6, 5.1.1, Exercise 13].

REFERENCES

- [1] Agnihotri and C. Woodward, *Eigenvalues of products of unitary matrices and quantum Schubert calculus*, Math. Research Letters **5** (1998), 817-836
- [2] A. Bertram, I. Ciocan-Fontanine and W. Fulton, *Quantum multiplication of Schur polynomials*, J. Algebra **219** (1999), 728-746
- [3] A. S. Buch, *Quantum Cohomology of the Grassmannians*, Preprint math.AG/0106268 (1. Jul 2001)

- [4] W. Fulton, *Young tableaux*, London Mathematical Society Student Texts: 35, Cambridge University Press, Cambridge, 1997
- [5] H. Hengelbrock, *An involution on the quantum cohomology ring of the Grassmannian*, Preprint math.AG/0205260 (24. May 2002)
- [6] D. E. Knuth, *The art of computer programming*, Vol. 3, Addison Wesley, (1973)
- [7] I.G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford mathematical monographs (1979)
- [8] K. Rietsch, *Quantum cohomology rings of Grassmannians and total positivity*, Preprint math.QA/0112022 (3. Dec 2001)
- [9] A. Postnikov *Affine approach to quantum Schubert calculus*, Preprint math.CO/0205165 (15. May 2002)
- [10] A. Postnikov *Symmetries of Gromov-Witten invariants* Advances in algebraic geometry motivated by physics, Contemp. Math., 276, (2000), 251-258
- [11] B. Siebert and G. Tian, *On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator*, Asian J. Math (1997), 679-695
- [12] R. P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, (1999)
- [13] J. Stembridge, *Computational Aspects of Root Systems, Coxeter Groups, and Weyl Characters*, Interactions of Combinatorics and Representation Theory, MSJ Memoirs **11** (2001), Math. Soc. Japan, (1-38)

Current address: Fakultät für Mathematik, Ruhr-Universität Bochum, 44803 Bochum

E-mail address: harald.hengelbrock@ruhr-uni-bochum.de